

Non-asymptotic confidence intervals for MCMC

Benjamin Gyori¹ and Daniel Paulin²

¹ *NUS Graduate School for Integrative Sciences and Engineering*
e-mail: bgyori@nus.edu.sg

² *Department of Mathematics*
e-mail: paulindani@gmail.com

National University of Singapore
21 Lower Kent Ridge Road, Singapore 119077, Republic of Singapore.

Abstract: Using concentration inequalities, we give non-asymptotic confidence intervals for estimates obtained by Markov chain Monte Carlo (MCMC) simulations, when using the approximation $\mathbb{E}_\pi f \approx (1/N) \cdot \sum_{i=1}^N f(X_i)$. We state results that are applicable on Markov chains on discrete as well as general state spaces. For reversible chains, we give an error bound depending explicitly on the spectral gap of the chain. In the non-reversible case, we formulate results using the chain's mixing time. We illustrate our results with simulations on lattice models in statistical physics as well as an example of Bayesian model averaging.

Keywords and phrases: Markov chain Monte Carlo, error bounds, non-asymptotic, confidence interval, concentration inequality, simulation.

AMS 2000 subject classifications: Primary 65C05, 60J10, 62M05; secondary 82B20, 68Q87, 68W20.

1. Introduction

The Monte Carlo method was invented by John von Neumann in the Los Alamos Laboratory, in 1947, for solving the problem of neutron diffusion in fissionable material, and thus helping Edward Teller to build the hydrogen bomb (see [Metropolis \(1987\)](#)).

The Monte Carlo method relies on independent samples from a probability distribution, to approximate an integral with respect to that distribution. Often, however, it is impossible or impractical to obtain such independent samples. One may still be able to construct a Markov chain with the target distribution as its stationary distribution. It is then possible to obtain a series of dependent samples by sampling from the Markov chain. This method is called Markov chain Monte Carlo (MCMC).

Let $X_1, X_2, \dots$, be a countable state, time homogeneous, ergodic Markov chain, taking values in Ω , and having stationary distribution π . Suppose that we are interested in computing $\mathbb{E}_\pi f$ for some $f : \Omega \rightarrow \mathbb{R}$. Then we usually make the approximation

$$\mathbb{E}_\pi f \approx \frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0}, \quad (1.1)$$

for some $t_0 \geq 0$ (“burn-in time”). For t_0 fixed, and $N \rightarrow \infty$, this average converges to $\mathbb{E}_\pi f$ by the ergodic theorem. However it is not clear how much this convergence is slowed down due to

the dependence of the samples. Consequently, an important question in practice is, how large should N be so that this approximation is correct to a certain level of precision? Practitioners often disregard this question, and keep silent about error bounds.

It is well known that the average in (1.1) tends to converge faster for fast mixing chains than for slow mixing ones. But until now, there have been few practically applicable results that relate the error in (1.1) to the mixing time and the spectral gap of the chain (for one such result, see [Lezaud \(1998a\)](#)). Most of the results in the literature are based on asymptotic convergence of the average to normal distribution. As we will see from our simulation results, such asymptotic bounds may underestimate the error for finite sample sizes. We will briefly review some of these results in Section 3.

Concentration inequalities can establish non-asymptotic error bounds for MCMC empirical averages of the form (1.1). For reversible chains, the speed of convergence is determined by the spectral gap when a sufficiently large burn-in time is chosen. In the non-reversible case the mixing time of the chain gives an upper bound on the speed of convergence. [Paulin \(2012a\)](#) establishes Hoeffding and Bernstein inequalities for both of these cases. In Section 6 of [Lezaud \(1998b\)](#), Bernstein inequalities are proven for Markov chains with general state space.

The purpose of this paper is to present these inequalities in a simple way, and show their applicability on simulation results of various models. We first look at simulations on lattice models in statistical physics, where the spectral gap and the mixing time are known. Then we present a case study of Bayesian inference, where the mixing properties of the chain are unknown. We have found that our bounds compare favorably with the existing asymptotic results.

1.1. Preliminary definitions

We define the mixing time of a time homogeneous chain as in Section 4.5 and 4.6 of [Levin, Peres and Wilmer \(2009\)](#). Let $X_1, X_2, X_3, \dots$ be a countable state, time homogeneous, irreducible, aperiodic Markov chain with transition matrix P , state space Ω , and stationary distribution π .

Let us denote

$$d(t) := \sup_{x \in \Omega} d_{TV}(P^t(x, \cdot), \pi),$$

where d_{TV} is the total variation distance. The mixing time of the chain is then defined as

$$t_{\text{mix}}(\epsilon) := \min\{t : d(t) \leq \epsilon\}$$

and

$$t_{\text{mix}} := t_{\text{mix}}(1/4).$$

If Ω is finite, we can write the eigenvalues of the transition matrix P as

$$1 = \lambda_1 > \lambda_2 \geq \dots \geq \lambda_{|\Omega|} > -1,$$

and we denote $\gamma := 1 - \lambda_2$ the *spectral gap*.

The mixing time and the spectral gap are related by the following inequality.

Lemma 1.1. *For reversible, irreducible, aperiodic chains in discrete time with finite state space Ω , denote*

$$t'_{\text{mix}} := \inf_{0 \leq \epsilon < 1/2} \frac{t_{\text{mix}}(\epsilon)}{\log\left(\frac{1}{2\epsilon}\right)} + 1, \quad (1.2)$$

for which

$$\frac{1}{\gamma} \leq t'_{\text{mix}} \leq \frac{t_{\text{mix}}}{\log(4)} + 1. \quad (1.3)$$

Proof. The first inequality follows from Theorem 12.4 of [Levin, Peres and Wilmer \(2009\)](#), for the second one, let $\epsilon = 4^{-n}$, then $t_{\text{mix}}(\epsilon) \leq nt_{\text{mix}}$, and $n \rightarrow \infty$ gives this bound. $\square$

For a random vector with distribution π , there are many ways to define a Markov chain that has π as stationary distribution.

Two of the most frequently used are the Metropolis-Hastings chain and the Gibbs sampler. Here we define the most frequently used variants of these (based on Chapter 3 of [Levin, Peres and Wilmer \(2009\)](#)).

Definition (Metropolis-Hastings chain). *Let Ω be any finite set, and Ψ an irreducible transition matrix. The Metropolis-Hastings chain modifies Ψ to obtain a chain with stationary distribution π .*

The transition matrix of the Metropolis-Hastings chain for a probability distribution π and symmetric transition matrix Ψ is defined as

$$P(x, y) = \begin{cases} \Psi(x, y) \min \left\{ 1, \frac{\pi(y)\Psi(y, x)}{\pi(x)\Psi(x, y)} \right\} & \text{if } y \neq x, \\ 1 - \sum_{z: z \neq x} \Psi(x, z) \min \left\{ 1, \frac{\pi(z)\Psi(z, x)}{\pi(x)\Psi(x, z)} \right\} & \text{if } y = x. \end{cases} \quad (1.4)$$

Remark 1.1. *In most of the practical situations $\pi(x) = h(x)/Z$, with Z being a normalization constant that is difficult to determine. A very important feature of the Metropolis-Hastings chain is that the transition probabilities only depend on π through the ratio $\frac{\pi(y)}{\pi(x)}$, which is independent of Z . The same holds true for the conditional probabilities in the case of the Gibbs sampler.*

The Gibbs sampler is a special case of the Metropolis-Hastings chain, when one can directly sample from the conditional distribution of each of the variables given the rest.

Definition (Gibbs sampler chain). *Assume that $\mathcal{S}$ is a finite set, $\mathcal{V}$ is a set of random variables, $\Omega = \mathcal{S}^{\mathcal{V}}$, and let π be a distribution on Ω . Then we define the Gibbs sampler chain as picking one variable in $\mathcal{V}$ uniformly at random, and resampling its value conditionally on the values on the rest of the variables.*

2. Results

In this section, we present concentration inequalities that give non-asymptotic bounds on the approximation (1.1). We state Hoeffding and Bernstein inequalities for both reversible and non-reversible chains.

2.1. Reversible chains

For reversible chains with countable state spaces, the following version of Hoeffding inequality holds (Theorem 1 of [León and Perron \(2004\)](#), the current form follows Theorem 2.7 of [Paulin \(2012a\)](#)):

Theorem 2.1 (Hoeffding inequality for reversible Markov chains). *Let $X = (X_1, \dots, X_N)$ be a time homogeneous, reversible, irreducible, aperiodic Markov chain taking values in some countable state space Ω , with stationary distribution π . Let λ_2 be the second largest eigenvalue of P ($\lambda_2 = 1 - \gamma$), t_{mix} the mixing time of the chain, and $f : \Omega \rightarrow [a, b]$. Let $t_0 \geq 0$ (“burn-in time”), and denote $Z := \frac{\sum_{i=t_0}^N f(X_i)}{N-t_0}$, and let $\lambda' = \max(0, \lambda_2)$. Then for any initial distribution and any $t \geq 0$,*

$$\begin{aligned} & \mathbb{P}[Z \geq \mathbb{E}_\pi f + t], \mathbb{P}[Z \leq \mathbb{E}_\pi f - t] \\ & \leq \exp\left(-2\frac{1-\lambda'}{1+\lambda'}(N-t_0)t^2/(b-a)^2\right) + \inf_{0 \leq \epsilon < 1} \epsilon^{\lfloor \frac{t_0}{t_{\text{mix}}(\epsilon)} \rfloor} \end{aligned} \quad (2.1)$$

Now we present a Bernstein-type result for countable state reversible chains (Corollary 2.10 of Paulin (2012a), which is based on the proof of Theorem 1.1 of Lezaud (1998a), see also Lezaud (1998b)):

Theorem 2.2 (Bernstein inequality for reversible Markov chains). *Let $X = (X_1, \dots, X_N)$ be a time homogeneous, reversible, irreducible, aperiodic Markov chain taking values in some countable state space Ω , with stationary distribution π , spectral gap γ , and mixing time t_{mix} . Suppose that $f : \Omega \rightarrow [-C, C]$, denote $V_f := \text{Var}_\pi(f) > 0$, and let*

$$C' := \sup_{x \in \Omega} |f(x) - \mathbb{E}_\pi f| \leq |E_\pi f| + C \leq 2C. \quad (2.2)$$

Let $t_0 \geq 0$ (“burn-in time”), and define $Z := \frac{\sum_{i=t_0+1}^N f(X_i)}{N-t_0}$, then for $t \geq 0$,

$$\begin{aligned} & \mathbb{P}[Z \geq \mathbb{E}_\pi f + t], \mathbb{P}[Z \leq \mathbb{E}_\pi f - t] \\ & \leq e^{\gamma/5} \exp\left[-\frac{(N-t_0)t^2\gamma}{4V_f(1+h(5tC'/V_f))}\right] + \inf_{0 \leq \epsilon < 1} \epsilon^{\lfloor \frac{t_0}{t_{\text{mix}}(\epsilon)} \rfloor} \end{aligned} \quad (2.3)$$

$$\leq e^{\gamma/5} \exp\left[-\frac{(N-t_0)t^2\gamma}{4V_f + 10C' \cdot t}\right] + \inf_{0 \leq \epsilon < 1} \epsilon^{\lfloor \frac{t_0}{t_{\text{mix}}(\epsilon)} \rfloor}, \quad (2.4)$$

with

$$h(x) := \frac{1}{2} \left(\sqrt{1+x} - (1-x/2) \right). \quad (2.5)$$

Define the asymptotic variance, σ^2 , as

$$\sigma^2 := \lim_{N \rightarrow \infty} \frac{1}{N} \text{Var}_\pi(f(X_1) + \dots + f(X_N)), \quad (2.6)$$

then the following bound holds:

$$\begin{aligned} & \mathbb{P}[Z \geq \mathbb{E}_\pi f + t], \mathbb{P}[Z \leq \mathbb{E}_\pi f - t] \leq \inf_{0 \leq \epsilon < 1} \epsilon^{\lfloor \frac{t_0}{t_{\text{mix}}(\epsilon)} \rfloor} + e^{\gamma/5} \cdot \\ & \cdot \exp\left[-(N-t_0) \cdot \left(\frac{\sqrt{\left(\sigma^2 + \frac{5}{\gamma}tC'\right)^2 + 4\sigma^2 K' t C'} - \left(\sigma^2 + \frac{5}{\gamma}tC'\right)}{2\sigma^2 K'} \right) \cdot \frac{t}{C'} \right], \end{aligned} \quad (2.7)$$

with $K' := \frac{10V_f}{\gamma^2\sigma^2} - \frac{5}{\gamma}$.

Remark 2.1. $\inf_{0 \leq \epsilon < 1} \epsilon^{\lfloor \frac{t_0}{t_{\text{mix}}(\epsilon)} \rfloor}$ can be further bounded by $4^{-\lfloor \frac{t_0}{t_{\text{mix}}} \rfloor}$. In many situations, Markov chains have a cutoff, which means that instead of geometrical decay, $t_{\text{mix}}(\epsilon)/t_{\text{mix}}(1-\epsilon) \rightarrow 1$ for any $0 < \epsilon < 1/2$ as the system size tends to infinity (see Figure 1 of [Lubetzky and Sly \(2009\)](#)). In such cases, choosing t_0 to be slightly larger (or a few times larger) than t_{mix} is sufficient.

2.2. Non-reversible chains

Most MCMC methods use reversible chains, in particular, the Metropolis-Hastings algorithm and the Gibbs sampler defined previously are reversible. However, using non-reversible chains can speed up the mixing time in some cases, for an example, see [Diaconis, Holmes and Neal \(2000\)](#). Therefore it is of interest to prove Hoeffding and Bernstein inequalities without assuming reversibility.

For later use, we define the following quantities (see also Proposition 1.2 of [Paulin \(2012a\)](#)):

$$t_{\text{mix}}^{\min} := \inf_{0 \leq \epsilon < 1} t_{\text{mix}}(\epsilon/2)/(1-\epsilon)^2 \leq 2.62t_{\text{mix}}, \quad (2.8)$$

$$t_{\text{mix}}^{\min'} := \inf_{0 \leq \epsilon < 1} t_{\text{mix}}(\epsilon/2)/(1-\sqrt{\epsilon})^2 \leq 4.43t_{\text{mix}}, \quad (2.9)$$

$$\eta_{\min}(t_0) := \inf_{0 \leq \epsilon < 1} \epsilon^{\lfloor \frac{t_0}{t_{\text{mix}}(\epsilon)} \rfloor} \cdot \frac{t_{\text{mix}}(\epsilon)}{1-\epsilon} \leq 4^{-\lfloor \frac{t_0}{t_{\text{mix}}} \rfloor} \cdot (4/3)t_{\text{mix}}. \quad (2.10)$$

For Markov chains with a cutoff, $t_{\text{mix}}^{\min} \approx t_{\text{mix}}^{\min'} \approx t_{\text{mix}}$.

The following two theorems are from [Paulin \(2012a\)](#) (Corollaries 2.4. and 2.9):

Theorem 2.3 (Hoeffding inequality for Markov chains). *First let $X = (X_1, \dots, X_N)$ be a time homogeneous Markov chain taking values in some countable space Ω . Suppose that $f : \Omega \rightarrow [a, b]$. Let $t_0 \geq 0$ (“burn-in time”), denote $Z := \frac{\sum_{i=t_0+1}^N f(X_i)}{N-t_0}$. Then for every $t \geq 0$,*

$$\begin{aligned} & \mathbb{P} \left(Z \geq \mathbb{E}_{\pi}(f) + \frac{\eta(t_0)(b-a)}{N-t_0} + t \right), \mathbb{P} \left(Z \leq \mathbb{E}_{\pi}(f) - \frac{\eta(t_0)(b-a)}{N-t_0} - t \right) \\ & \leq \exp \left(\frac{-2(N-t_0)t^2}{(b-a)^2 t_{\text{mix}}^{\min}} \right) \leq \exp \left(\frac{-(N-t_0)t^2}{1.31(b-a)^2 t_{\text{mix}}} \right). \end{aligned} \quad (2.11)$$

Remark 2.2. We give a short direct proof for this result in the Appendix.

Theorem 2.4 (Bernstein inequality for Markov chains). *Let $X = (X_1, \dots, X_N)$ be a homogeneous Markov chain taking values in some countable space Ω , with stationary distribution π . Let $f : \Omega \rightarrow [-C, C]$, let $t_0 \geq 0$ (“burn-in time”), and denote $Z := \frac{\sum_{i=t_0+1}^N f(X_i)}{N-t_0}$. Let C' be defined as in (2.2). Denote*

$$V := \sum_{i=t_0+1}^N \mathbb{E} (f(X_i) - \mathbb{E}_{\pi} f)^2. \quad (2.12)$$

Then for every $t \geq 0$,

$$\begin{aligned} & \mathbb{P} \left(Z \geq \mathbb{E}_{\pi}(f) + \frac{\eta(t_0)C'}{N-t_0} + t \right), \mathbb{P} \left(Z \leq \mathbb{E}_{\pi}(f) - \frac{\eta(t_0)C'}{N-t_0} - t \right) \\ & \leq \exp \left(\frac{-t^2(N-t_0)^2}{t_{\text{mix}}^{\min'}(8V + 4\sqrt{2}(N-t_0)C't)} \right) \\ & \leq \exp \left(\frac{-t^2(N-t_0)^2}{t_{\text{mix}}(35.5V + 25.1(N-t_0)C't)} \right). \end{aligned} \quad (2.13)$$

Remark 2.3. This form of Theorem 2.2 and 2.4 follows from Corollary 2.11, and Corollary 2.10 of Paulin (2012a) by the substitution $f \rightarrow f - \mathbb{E}_\pi f$. For another Bernstein inequality, which uses the pseudo spectral gap of the chain, instead of the mixing time, see Theorem 2.5 of Paulin (2012a).

2.3. Chains with general state space

Lezaud (1998b) also proves Bernstein inequalities for chains on general state spaces. However, some quantities appearing in these formulas are hard to interpret in practice. We now state two such results and explain how they can be employed in practice.

The following result is a Bernstein inequality for reversible chains on general state spaces, adapted from Lezaud (1998b), page 98 (here we have reformulated it to work for functions that are not necessarily centered, taking values in $[-C, C]$):

Theorem 2.5 (Theorem 1.1, Chapter 6 of Lezaud (1998b)). *Let $P(x, dy)$ be a Markov kernel over a probability space $(\Omega, \mathcal{F}, \pi)$, with π being the unique stationary distribution of P . Let $X_1, X_2, \dots$ be a Markov chain with this kernel. Let $f : \Omega \rightarrow \mathbb{R}$ be a bounded function, with $\|f\|_\infty \leq C$ and $V_f := \text{Var}_\pi(f) > 0$. Let C' be as in (2.2). For a distribution p on Ω that has a density φ relative to π , denote $N_p := (\mathbb{E}_\pi \varphi^2)^{1/2}$. Suppose that the linear operator P defined by $P(x, dy)$ extends to $L^2(\pi)$, and it is self-adjoint. Denote the initial distribution of the chain by q , and let $q(k) := P^k q$.*

Let $t_0 \geq 0$ (“burn-in time”) and denote $Z := \frac{\sum_{i=t_0+1}^N f(X_i)}{N-t_0}$. Then for every $t \geq 0$,

$$\begin{aligned} & \mathbb{P}_q [Z \geq \mathbb{E}_\pi f + t], \mathbb{P}_q [Z \leq \mathbb{E}_\pi f - t] \\ & \leq N_{q(t_0)} e^{\gamma/5} \exp \left[-\frac{(N-t_0)t^2\gamma}{4V_f(1+h(5tC'/V_f))} \right] \end{aligned} \quad (2.14)$$

$$\leq N_{q(t_0)} e^{\gamma/5} \exp \left[-\frac{(N-t_0)t^2\gamma}{4V_f + 10C't} \right], \quad (2.15)$$

where γ is the spectral gap of the chain and $h(x)$ is defined as in (2.5).

In practice, we often do not know how to estimate N_q . However, it is clear that for ergodic chains, $q(t_0) = P^{t_0} q \rightarrow \pi$ as $t_0 \rightarrow \infty$, and thus $N_{q(t_0)} \rightarrow 1$ in most situations of practical interest. Therefore, for reversible chains, for an arbitrarily initial distribution, if we choose t_0 sufficiently large, then roughly same bounds hold as in the countable state space case, see (2.4).

The following result is a Bernstein inequality for non-reversible Markov chains in general state space. It is an improvement on Theorem 1.5 on page 99 of Lezaud (1998b), similarly as it is done in Theorem 2.8 of Paulin (2012a):

Theorem 2.6. *Let $P(x, dy)$ be a Markov kernel over a probability space $(\Omega, \mathcal{F}, \pi)$, with π being the unique stationary distribution of P . Let $X_1, X_2, \dots$ be a Markov chain with this kernel. Let $f : \Omega \rightarrow \mathbb{R}$ be a bounded function, with $\|f\|_\infty \leq C$ and $V_f := \text{Var}_\pi(f) > 0$. Let C' be as in (2.2). For a distribution p on Ω that has a density φ relative to π , denote $N_p := (\mathbb{E}_\pi \varphi^2)^{1/2}$. Suppose that the linear operator P defined by $P(x, dy)$ extends to $L^2(\pi)$, and it is self-adjoint. Denote the initial distribution of the chain by q , and let $q(k) := P^k q$. Let $\gamma((P^*)^k P^k)$ be the spectral gap of the self-adjoint linear operator $(P^*)^k P^k$, and define the pseudo spectral gap as*

$$\gamma_{\text{ps}} := \sup_{k \geq 1} \frac{\gamma((P^*)^k P^k)}{k}. \quad (2.16)$$

Let $t_0 \geq 0$ (“burn-in time”) and denote $Z := \frac{\sum_{i=t_0+1}^N f(X_i)}{N-t_0}$. Then for every $t \geq 0$,

$$\begin{aligned} & \mathbb{P}_q [Z \geq \mathbb{E}_\pi f + t], \mathbb{P}_q [Z \leq \mathbb{E}_\pi f - t] \\ & \leq N_{q(t_0)} \exp \left[-\frac{(N-t_0)t^2\gamma_{\text{ps}}}{8V_f(1+h(5tC'/V_f))} \right] \end{aligned} \quad (2.17)$$

$$\leq N_{q(t_0)} \exp \left[-\frac{(N-t_0)t^2\gamma_{\text{ps}}}{8V_f + 20C't} \right], \quad (2.18)$$

with $h(x)$ defined as in the previous theorem.

Remark 2.4. The proof of this is left to the reader as an exercise.

Again, as in (2.5), $N_{q(t_0)} \rightarrow 1$ as $t_0 \rightarrow \infty$ in most practical cases. It is important to note that on general state spaces, the mixing time based on the total variation distance can not be used. However, the pseudo spectral gap can be used as an alternative to the mixing time in the non-reversible case.

2.4. Subsampling

$Z := \frac{\sum_{i=t_0+1}^N f(X_i)}{N-t_0}$ is not the only possible way to approximate $\mathbb{E}_\pi f$. We may decide to only average in every m th step (typically, we choose $m = 1/\gamma$ for reversible chains, or $m = t_{\text{mix}}$ for non-reversible chains). Assume, without loss of generality, that

$$N = nm \text{ and } t_0 = t'_0 m. \quad (2.19)$$

Denote $X'_1 := X_m, X'_2 := X_{2m}, \dots, X'_n := X_{n \cdot m}$, and

$$Z' := \frac{\sum_{i=t'_0+1}^n f(X'_i)}{n-t'_0}. \quad (2.20)$$

Then $X'_1, \dots, X'_n$ is a Markov chain, which is reversible if the original chain was reversible. In this case, choose m to be odd, then the new transition matrix P^m will have second largest eigenvalue λ^m (where λ denotes the second largest eigenvalue of P), and thus its spectral gap is $\gamma' = 1 - (1 - \gamma)^m$. Let m_γ denote the smallest odd number greater or equal to $1/\gamma$, then with the choice $m := m_\gamma$, we have $\gamma' = 1 - (1 - \gamma)^m \geq \frac{e-1}{e}$ (this also holds in case $\gamma > 1$). Similarly, for non-reversible chains, with the choice $m = t_{\text{mix}}$, $X'_1, \dots, X'_n$ will have mixing time 1.

Therefore, with these choices, the reader can see that almost the same concentration inequalities hold for Z' as for Z (by applying our theorems on Z'). The advantage of this approach is that one only needs to compute f in every $1/\gamma$ -th (or t_{mix} -th) step, which may result in considerable savings if f is expensive to evaluate.

3. Comparison with previous results

In this section we give a brief review of some widely used MCMC convergence diagnostics and error estimation methods. For a more comprehensive overview of available techniques, we refer the reader to [Cowles and Carlin \(1996\)](#), [Liu \(2008\)](#) and [Diaconis \(2011\)](#).

3.1. Convergence diagnostics

The most frequently used convergence analysis method, the Gelman-Rubin diagnostic, was introduced in Gelman and Rubin (1992) (this was further refined in Brooks and Gelman (1998), see also Gelman et al. (2004)).

Gelman and Rubin (1992) propose a multiple sequence convergence assessment method: first, m parallel chains are run for $2n$ steps. Then the first n terms of each chain are thrown away. Finally, B and W (the *between*, and *within sequence variations*) are computed for each estimated function f of interest:

$$B := \frac{n}{m-1} \sum_{j=1}^m (\hat{E}^{(j)} - \bar{\hat{E}})^2 \text{ where } \hat{E}^{(j)} := \sum_{i=1}^n f(X_i^{(j)})/n, \bar{\hat{E}} := \sum_{j=1}^m \hat{E}^{(j)}/m. \quad (3.1)$$

$$W := \frac{1}{m} \sum_{j=1}^m V^{(j)} \text{ where } V^{(j)} := \sum_{i=1}^n \left(f(X_i^{(j)}) - \hat{E}^{(j)} \right)^2 / (n-1) \quad (3.2)$$

From these, one computes the *potential scale reduction*, $\hat{R}$, as a function of B and W .

$$\hat{R} := \sqrt{\frac{((n-1)/n)W + (1/n)B}{W}}. \quad (3.3)$$

Furthermore, one can estimate the *effective number of independent samples*, n_{eff} , as

$$n_{\text{eff}} := mn \frac{((n-1)/n)W + (1/n)B}{B}. \quad (3.4)$$

This estimate corresponds to the aggregate number of “independent” samples from the m parallel runs. The chain is assumed to have reached convergence when $\hat{R}$ is sufficiently close to 1. The authors recommend $\hat{R} \leq 1.1$ as a threshold adequate in most situations (for more details, see pages 294-298 of Gelman et al. (2004)). The main strength of this approach is that it is easy to implement, and is available in most statistical packages. However, it does not offer a quantitative bound on the error of the estimate $\frac{1}{n} \sum_{i=1}^n f(X_i) - \mathbb{E}_\pi f$.

3.2. Error estimation by the central limit theorem

A central limit theorem (CLT) for Markov chains was introduced in Kipnis and Varadhan (1986):

Theorem 3.1. *Let $(X_i)_{i \geq 1}$ be a stationary, irreducible, reversible Markov chain taking values in some general state space Ω , with stationary distribution π . Let $f : \Omega \rightarrow \mathbb{R}$, $Z_n := \frac{1}{n} \sum_{i=1}^n f(X_i)$, and $\gamma_t := \text{Cov}_\pi(f(X_i), f(X_{i+t}))$. Then*

$$n \text{Var}(Z_n) \rightarrow \sigma^2 := \sum_{t=-\infty}^{\infty} \gamma_t \text{ almost surely.} \quad (3.5)$$

If σ^2 is finite, then

$$\sqrt{n}(Z_n - \mathbb{E}_\pi f) \Rightarrow N(0, \sigma^2). \quad (3.6)$$

Remark 3.1. This form of the theorem appears in [Geyer \(1992\)](#). For a more precise statement, see Theorem 17.0.1 of [Meyn and Tweedie \(2009\)](#).

The conditions of the above theorem are satisfied by a large class of chains, for instance, both Gibbs and Metropolis-Hastings sampling (as defined in Section 1.1) are reversible.

To make use of the limiting distribution $N(0, \sigma^2)$, [Geyer \(1992\)](#) proposes several estimators of σ^2 . Firstly, the *lagged autocovariance* γ_t is estimated by the *empirical autocovariance*

$$\hat{\gamma}_{n,t} = \hat{\gamma}_{n,-t} := \frac{1}{n} \sum_{i=1}^{n-t} [f(X_i) - Z_n] [f(X_{i+t}) - Z_n]. \quad (3.7)$$

Define $\Gamma_{n,m} := \hat{\gamma}_{n,2m} + \hat{\gamma}_{n,2m+1}$. The *initial positive sequence estimator* is defined as

$$\hat{\sigma}_{pos,n}^2 := \hat{\gamma}_{n,0} + 2 \sum_{i=1}^{2m+1} \hat{\gamma}_{n,i} = -\hat{\gamma}_{n,0} + 2 \sum_{i=0}^m \hat{\Gamma}_{n,i}, \quad (3.8)$$

where m is chosen to be the largest integer such that $\hat{\Gamma}_{n,i} > 0$ for $1 \leq i \leq m$. The authors also define the *initial monotone sequence estimator* $\hat{\sigma}_{mon,n}^2$ by replacing $\hat{\Gamma}_{n,i}$ by $\min_{j \leq i} \hat{\Gamma}_{n,j}$, and the *initial convex sequence estimator* $\hat{\sigma}_{conv,n}^2$ by taking the greatest convex minorant.

For all of these three estimators, [Geyer \(1992\)](#) proves that for almost all sample paths,

$$\liminf_{n \rightarrow \infty} \hat{\sigma}_n^2 \geq \sigma^2,$$

i.e. the variance σ^2 is asymptotically overestimated. Therefore asymptotically conservative confidence intervals can be obtained by using the quantiles of $N(0, \hat{\sigma}_n^2)$ with any of the three estimators.

The main advantage of the above method is that it applies for general state space reversible chains, with any square integrable f . The disadvantage is that it is only proven to work asymptotically, and we often do not know how well the Kipnis-Varadhan CLT approximates the normal distribution. Even in the independent case, by the Berry-Esseen bound, there is an error of order $\frac{1}{\sqrt{N}}$ in Kolmogorov distance. For Markov chains, this can be higher, especially when the mixing is slow.

The advantage of our method is that it works non-asymptotically, and thus we can get more reliable error estimates when the mixing time and spectral gap of the chain can be bounded.

3.3. Error bounds via Ricci curvature

Ricci curvature for proving concentration of empirical averages of functions of Markov chains was introduced in [Ollivier \(2009\)](#), and further developed in [Joulin and Ollivier \(2010\)](#). Here, we will give a simple exposition of this method and compare it with our results.

Let (Ω, d) be a Polish space (metric, complete and separable) and define a Markov chain $X_1, X_2, \dots$ on this space with unique stationary distribution π . We denote the associated transition kernel $(P_x)_{x \in \Omega}$ so that P_x is a probability measure on Ω and $P_x(dy)$ is the transition probability from x to y . The N -step transition kernel will be denoted by P_x^N . The Ricci curvature is defined in terms of the Wasserstein distance. The *Wasserstein distance* of two measures, μ_1 and μ_2 on (Ω, d) is defined as:

$$W_1(\mu_1, \mu_2) := \inf_{\Pi[X \sim \mu_1, Y \sim \mu_2]} \mathbb{E}_{\Pi} (1[X \neq Y]), \quad (3.9)$$

ie. the infimum is taken over all couplings of μ_1 and μ_2 .

We say that a Markov chain has positive *Ricci curvature* κ if it satisfies the following assumption:

Assumption 3.1. *There exists $\kappa > 0$ such that*

$$W_1(P_x, P_y) \leq (1 - \kappa)d(x, y), \quad (3.10)$$

for any $x, y \in \Omega$.

Define the *eccentricity* E at point $x \in \Omega$ as

$$E(x) := \mathbb{E}_{Y \sim \pi} [d(x, Y)]. \quad (3.11)$$

Let $\sigma(x)^2$ denote the coarse diffusion constant defined as

$$\sigma(x)^2 := \frac{1}{2} \mathbb{E}_{\pi_i} [d(Y, Z)^2], \quad (3.12)$$

where π_i is an independent coupling between the random variables $Y \sim P_x, Z \sim P_x$.

The local dimension $n_x, x \in \Omega$ and is defined as

$$n_x := \inf_{f: \Omega \rightarrow \mathbb{R}, \|f\|_{\text{Lip}} \leq 1} \frac{\mathbb{E}_{\pi_i} [d(Y, Z)^2]}{\mathbb{E}_{\pi_i} [|f(Y) - f(Z)|^2]} \geq 1, \quad (3.13)$$

where $\|f\|_{\text{Lip}} = \sup_{x \neq y} (|f(x) - f(y)|/d(x, y))$.

We define the granularity of the Markov chain as

$$\sigma_\infty := \frac{1}{2} \sup_{x \in \Omega} \text{diam Supp } P_x, \quad (3.14)$$

where Supp refers to support.

Theorem 3.2 (Theorem 4 of [Joulin and Ollivier \(2010\)](#)). *Let $Z = \frac{\sum_{i=t_0+1}^N f(X_i)}{N-t_0}$ and define*

$$V^2 = \frac{1}{\kappa(N-t_0)} \left(1 + \frac{t_0}{N-t_0} \right) \sup_{x \in \Omega} \frac{\sigma^2(x)}{n_x \kappa} \quad (3.15)$$

then the following concentration inequality holds ($\mathbb{P}_x$ and $\mathbb{E}_x$ refers to initial point $X_1 = x$)

$$\begin{aligned} & \mathbb{P}_x(Z \geq \mathbb{E}_x(Z) + t), \mathbb{P}_x(Z \leq \mathbb{E}_x(Z) - t) \\ & \leq \begin{cases} 2 \exp(-t^2/(16V^2\|f\|_{\text{Lip}}^2)) & \text{if } t \geq t_{\max} \\ 2 \exp(-\kappa(N-t_0)t/(12\sigma_\infty\|f\|_{\text{Lip}})) & \text{if } t \geq t_{\max}, \end{cases} \end{aligned} \quad (3.16)$$

the boundary of the Gaussian window is $t_{\max} := 4V^2\kappa(N-t_0)\|f\|_{\text{Lip}}/(3\sigma_\infty)$. The bias can be bounded as in Proposition 1 of [Joulin and Ollivier \(2010\)](#):

$$|\mathbb{E}_x Z - \mathbb{E}_\pi Z| \leq \frac{(1-\kappa)^{t_0+1}}{\kappa(N-t_0)} E(x) \|f\|_{\text{Lip}}. \quad (3.17)$$

We later show how κ can be bounded from below on spin lattice models, see Section 5.

The above result depends on the distance d and $\|f\|_{\text{Lip}}$ with respect to this distance. This can be advantageous for some general state space models but in discrete state space, its sensitivity may ruin the tightness of the bound.

Our results employ well established quantities of the chain – the spectral gap in the reversible case and the mixing time in the non-reversible case – and depend only on the boundedness of f , as $\|f\|_\infty$. We also provide methods for estimating all parameters appearing in our inequalities, which makes them easier to apply in practice.

3.4. Estimation of the mean square error

Non-asymptotic results also exist for the mean square error of the MCMC estimate. Generally speaking, these results work even for unbounded functions f that have a finite variance (with respect to π). However, their main weakness is that one can only deduce quadratic decay through the Chebyshev inequality instead of exponential or Gaussian decay as with concentration inequalities.

We present a simple example of the Chebyshev inequality for finite state, reversible Markov chains:

Theorem 3.3 (Theorem 12.19. of [Levin, Peres and Wilmer \(2009\)](#)). *Let $(X_t)_{t \geq 1}$ be a finite state, irreducible, aperiodic, reversible Markov chain, with stationary distribution π , and spectral gap γ . If $t_0 \geq t_{\text{mix}}(\epsilon/2)$ and $N \geq \frac{1}{\gamma} \cdot [4 \text{Var}_\pi(f)/(\eta^2 \epsilon)] + t_0$, denote $Z := \frac{\sum_{i=t_0}^N f(X_i)}{N-t_0}$, then (for any initial distribution)*

$$\mathbb{P}\{|Z - \mathbb{E}_\pi f| \geq \eta\} \leq \epsilon. \quad (3.18)$$

Remark 3.2. $\text{Var}_\pi(f)$ can be estimated as in (4.1).

Further non-asymptotic bounds have also been proposed based on mean square error. [Rudolf \(2011\)](#) gives a similar bound on the mean square error using the spectral gap γ and $\|f\|_p$ for $p \geq 2$. [Latuszynski, Miasojedow and Niemiro \(2011\)](#) gives an asymptotically sharp bound on the mean square error as a function of the asymptotic variance σ^2 . The bounds are valid on general state spaces without assuming reversibility.

4. Estimation of parameters

The main difficulty we encounter when applying Theorem 2.2 is that, in general, we do not know $V_f = \text{Var}_\pi(f)$ and σ^2 (see (2.6)). Similarly, in Theorem 2.4, we usually do not know V (see (2.12)). In many cases, the spectral gap γ and mixing time t_{mix} are also unknown.

4.1. Estimation of the variance

From the definitions, it is easy to see that we can estimate V_f as

$$\hat{V}_f := \left(\frac{1}{N-t_0} \sum_{i=t_0+1}^N f^2(X_i) \right) - \left(\frac{1}{N-t_0} \sum_{i=t_0+1}^N f(X_i) \right)^2. \quad (4.1)$$

Similarly, we can estimate V as

$$\begin{aligned} \hat{V} &:= \sum_{i=t_0+1}^N \left(f(X_i) - \frac{1}{N-t_0} \left(\sum_{i=t_0+1}^N f(X_i) \right) \right)^2 \\ &= \sum_{i=t_0+1}^N f^2(X_i) - \frac{1}{N-t_0} \left(\sum_{i=t_0+1}^N f(X_i) \right)^2 = (N-t_0) \hat{V}_f. \end{aligned} \quad (4.2)$$

Our next proposition gives a bound on the upper tails of $V_f - \hat{V}_f$ and $V - \hat{V}$:

Proposition 4.1. *Let $\eta_{\min}(t_0)$ be as in (2.10). Then we have, for any $T \geq 0$,*

$$\mathbb{P} \left(V_f - \hat{V}_f \geq \frac{5\eta_{\min}(t_0)C^2}{N - t_0} + T \right) \leq 3 \exp \left(\frac{-2(N - t_0)T^2}{9C^4 t_{\min}^{\min}} \right), \quad (4.3)$$

$$\mathbb{P} \left(V - \hat{V} \geq 10\eta_{\min}(t_0)C^2 + T \right) \leq 3 \exp \left(\frac{-2T^2}{9(N - t_0)C^4 t_{\min}^{\min}} \right). \quad (4.4)$$

Remark 4.1. *In practice, in most cases, we will use $\hat{V}_f$ and $\hat{V}$ directly, and this proposition ensures that for sufficiently large N , the mistake is negligible.*

We will use the monotone sequence estimator of Geyer (1992) to estimate σ^2 (see Section 3). We denote this estimate by $\hat{\sigma}^2 := \hat{\sigma}_{\text{mon}, N-t_0}^2$.

4.2. Estimation of the spectral gap and the mixing time

Precise estimation of the spectral gap and the mixing time from the realizations $f(X_1), \dots, f(X_N)$ is not possible, since it is a property of the Markov chain $X_1, \dots, X_N$ itself, and by applying the function f , we lose information. Nevertheless, in practice, we have found that the simple estimate in (4.6) works well. We now give a brief justification of this approach.

For reversible chains with state space Ω , transition matrix P , and stationary distribution π , define $l^2(\pi)$ as the Hilbert space of real valued functions on Ω , with scalar product

$$\langle f, g \rangle := \sum_{x \in \Omega} f(x)g(x)\pi(x).$$

Let $\{\varphi_i\}_{i \geq 1}$ be an orthonormal basis made of the eigenvectors of P , corresponding to eigenvalues $(\lambda_i)_{i \geq 1}$. Then the largest eigenvalue $\lambda_1 = 1$, and $\varphi_1 = 1$, and obviously $\lambda_2 = 1 - \gamma$.

By Proposition 1.5 on page 48 of Lezaud (1998b), we have

$$\sigma^2 = \sum_{i \geq 2} \langle f, \varphi_i \rangle^2 \frac{1 + \lambda_i}{1 - \lambda_i} \leq \frac{2V_f}{\gamma}, \quad (4.5)$$

thus $\gamma \leq 2V_f/\sigma^2$ for any function f . With the choice $f = \varphi_1$, we have $\sigma^2 = (2 - \gamma)/\gamma$, and $V_f = 1$, thus $2V_f/\sigma^2 = \gamma \cdot 2/(2 - \gamma)$, which is indeed very close to γ (in practice, $\gamma \ll 1$).

For this reason, if we are only equipped with the values of a single function $f: f(X_1), \dots, f(X_N)$, then we propose the estimate

$$\hat{\gamma} := \frac{2\hat{V}_f}{\hat{\sigma}^2}. \quad (4.6)$$

In case we have the values $f(X_1), \dots, f(X_N)$ for several functions $f_1, f_2, \dots, f_k$, then denote the corresponding estimates of $\hat{\sigma}^2$ and $\hat{V}_f$ by $\hat{\sigma}_1^2, \dots, \hat{\sigma}_k^2$ and $\hat{V}_{f_1}, \dots, \hat{V}_{f_k}$. We estimate γ as

$$\hat{\gamma} := \min_{1 \leq i \leq k} \frac{2\hat{V}_{f_i}}{\hat{\sigma}_i^2}. \quad (4.7)$$

A popular method for assessing convergence of the chain has been proposed by Gelman and Rubin (1992). We will make use of this method to indirectly estimate the mixing time. Having obtained an estimate of n_{eff} with m parallel chains taking n steps (see Section 3), we can obtain

an estimate of the mixing time. Since n_{eff} corresponds to the aggregate number of “independent” samples from m runs, n_{eff}/m is the average number of such samples per run. Following our argument on subsampling in Section 2.4, we propose the following estimate for the mixing time:

$$\hat{t}_{\text{mix}} := \frac{nm}{n_{\text{eff}}}. \quad (4.8)$$

With $\hat{\gamma}$ estimated as in (4.6) and $\hat{t}_{\text{mix}}$ as in (4.8), the Bernstein inequality for reversible chains becomes

$$\begin{aligned} & \mathbb{P}[Z \geq \mathbb{E}_{\pi} f + t], \mathbb{P}[Z \leq \mathbb{E}_{\pi} f - t] \\ & \leq \exp\left(\frac{2\hat{V}_f}{5\hat{\sigma}^2}\right) \cdot \exp\left[-\frac{(N - t_0)t^2}{2\hat{\sigma}^2 + 5\frac{\hat{\sigma}^2 C'}{\hat{V}_f} \cdot t}\right] + 4^{-\lfloor \frac{t_0}{\hat{t}_{\text{mix}}} \rfloor}, \end{aligned} \quad (4.9)$$

with C' defined as in (2.2). Assuming that our estimate of t_{mix} in (4.8) is the correct order of magnitude, we can choose t_0 large enough for the second term to become negligible.

4.3. Advice to practitioners

When applying MCMC methods in practice, the mixing time and spectral gap of the chain is often not known. Using estimates for the relevant parameters as described above, we recommend the following procedure (for reversible chains):

1. Estimate t_{mix} using (4.8) (if too close to n , repeat with larger n).
2. Run two parallel chains for $k\hat{t}_{\text{mix}}$ iterations. We recommend using $k = 1000$ based on Proposition 4.1. Set $t_0 = 10\hat{t}_{\text{mix}}$. Estimate $\hat{\sigma}^2$ and $\hat{V}_f$ for both chains (using the monotone sequence estimator $\hat{\sigma}_{\text{mon}, N-t_0}^2$, and (4.1)). If, for these 2 chains, the estimated values are far away, then increase the number of iterations.
3. Use (4.9) with the estimated values of $\hat{\sigma}$ and $\hat{V}_f$ to obtain the necessary number of steps for a given precision, and make the final simulation with this amount of steps.

5. Simulations

In the following, we present simulation results to demonstrate the applicability of the introduced error bounds. We are interested in the empirical tail probabilities of estimates, obtained from multiple runs of MCMC simulations. In particular, we will estimate logarithms of tail probabilities of the following form:

$$\log\left(\mathbb{P}\left(\frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0} \geq \mathbb{E}_{\pi} f + t\right)\right). \quad (5.1)$$

We simulate m parallel chains and denote the sequence of states of the j th chain ($1 \leq j \leq m$) by $X_1^{(j)}, \dots, X_N^{(j)}$. Then the empirical average obtained by the j th chain can be written as

$$\hat{E}^{(j)} := \frac{\sum_{i=t_0+1}^N f(X_i^{(j)})}{N - t_0}, \quad (5.2)$$

and denote

$$\widehat{E} := \frac{1}{m} \sum_{j=1}^m \widehat{E}^{(j)}. \quad (5.3)$$

Define the *mean-shifted empirical distribution* of these estimates as

$$\widehat{F}(t) := \frac{1}{m} \sum_{j=1}^m \mathbb{I}[\widehat{E}^{(j)} - \widehat{E} \leq t], \quad (5.4)$$

and let

$$\widehat{L}(t) := \begin{cases} \log(\widehat{F}(t)) & \text{for } t < 0, \text{ and} \\ \log(1 - \widehat{F}(t)) & \text{for } t \geq 0, \end{cases} \quad (5.5)$$

thus $\widehat{L}(t)$ is an estimate of the log tails in (5.1).

5.1. Lattice models in statistical physics

We first consider simulations on the Curie-Weiss model and the Ising model (1D and 2D). These models and their variants are widely studied in the context of MCMC simulations, and for some special cases, the mixing time and spectral gap of the chain are known.

5.1.1. Definition of models

Let us assume that $\sigma := (\sigma_1, \dots, \sigma_{n_s})$ are spins taking values 1 or -1 , and distributed according to the probability distribution of the model, which is of the form (for some $\beta > 0$)

$$\mathbb{P}(\sigma_1, \dots, \sigma_{n_s}) = \frac{\exp(H_{\beta,h}(\sigma))}{Z}, \quad (5.6)$$

where $H_{\beta,h}(\sigma)$ is the energy function, β is the inverse temperature, h corresponds to the external field, and $Z = \sum_{\sigma} \exp(H_{\beta,h}(\sigma))$ is the partition function.

In the case of the Curie-Weiss model, we define the energy function as

$$H_{\beta,h}(\sigma) = H_{\beta,h}^{CW}(\sigma) := \frac{\beta}{n_s} \sum_{1 \leq i < j \leq n_s} \sigma_i \sigma_j + h \sum_{i=1}^{n_s} \sigma_i. \quad (5.7)$$

In the case of Ising model on a graph $G = (V, E)$, we say that $i \sim j$ if there is an edge between i and j in G , and define H as

$$H_{\beta,h}(\sigma) = H_{\beta,h}^I(\sigma) := \beta \sum_{i \sim j} \sigma_i \sigma_j + h \sum_{i=1}^{n_s} \sigma_i. \quad (5.8)$$

In the 1-dimensional Ising model, G consists of the edges $(i, i+1)$ for $1 \leq i \leq n_s - 1$, while in the 2 dimensional case, G consists of the edges on a square lattice. We use periodic boundary conditions so that each spin is connected with the same number of other spins.

In practice it is impossible to obtain independent samples from (5.6). Therefore one typically designs a Markov chain which has (5.6) as its stationary distribution. In the following section we present analytic estimates of the properties of one such typically used Markov chain, the Glauber dynamics (a Gibbs sampler chain on the lattice model, as defined in Section 1.1).

5.1.2. Spectral gap and mixing time

In the case of the Curie-Weiss model, at high temperature, with no external field, ($\beta < 1$ and $h = 0$), Theorem 1 of [Ding, Lubetzky and Peres \(2009\)](#) shows that for the Glauber dynamics, the spectral gap and mixing time are

$$\gamma^{CW} = \frac{(1 + o(1))(1 - \beta)}{n_s}, \quad t_{\text{mix}}^{CW} = \frac{1}{2} \frac{n_s \log((1 - \beta)^2 n_s)}{1 - \beta} + O\left(\frac{n_s}{1 - \beta}\right),$$

so we will use the following approximate values:

$$\hat{\gamma}^{CW} := \frac{1 - \beta}{n_s}, \quad \hat{t}_{\text{mix}}^{CW} := \frac{1}{2} \frac{n_s \log((1 - \beta)^2 n_s)}{1 - \beta}. \quad (5.9)$$

In the high temperature case ($\beta < 1$), but with $h \neq 0$, we can still apply Theorem 3 of [Bubley et al. \(1997\)](#) to show that the mixing time satisfies

$$t_{\text{mix}}(\epsilon) \leq \lceil n_s \log(n_s/\epsilon)/(1 - \beta) \rceil, \text{ and thus } t_{\text{mix}} \leq \lceil n_s \log(4n_s)/(1 - \beta) \rceil,$$

therefore we expect (5.9) to be good approximation in this case as well. [Ding, Lubetzky and Peres \(2009\)](#) also shows that for the critical ($\beta = 1$, $h = 0$) case, the mixing time is $O(n_s^{3/2})$. For low temperature ($\beta > 1$, $h = 0$), the mixing time is exponential in n_s . The inverse spectral gap, $1/\gamma$, has the same order as the mixing time for both the critical and the low temperature case respectively.

In the case of d dimensional Ising model with periodic boundaries (i.e. n_s^d spins in total), [Lubetzky and Sly \(2009\)](#) shows that for the continuous time Glauber dynamics (i.e. Glauber dynamics, with i.i.d rate-one Poisson clocks on each spin), the mixing time satisfies

$$t_{\text{mix}}^{d,Gl} = \frac{d}{2\gamma_{\infty}^{d,Gl}} \log(n_s) + O(\log(\log(n_s))), \quad (5.10)$$

where $\gamma_{\infty}^{d,Gl}$ is the spectral gap of the Glauber dynamics on the d dimensional infinite lattice. The same result holds for the Metropolis algorithm, with $\gamma_{\infty}^{d,Gl}$ replaced by $\gamma_{\infty}^{d,Metropolis}$.

The spectral gap of the finite model satisfies, by Theorem 4 of [Lubetzky and Sly \(2009\)](#): under some weak conditions (strong spatial mixing) on β and h , $|\gamma_{n_s}^{d,Gl} - \gamma_{\infty}^{d,Gl}| \leq n_s^{-1/2+o(1)}$.

For the 1-dimensional case with $h = 0$, strong spatial mixing holds, and $\gamma_{\infty}^{1,Gl} = \gamma^{1,Gl}(n_s) = 1 - \tanh(2\beta)$ (independently of n_s). This means that for $h = 0$,

$$\begin{aligned} t_{\text{mix}}^{1,Gl} &= \frac{1}{2\gamma_{\infty}^{1,Gl}} \log(n_s) + O(\log(\log(n_s))) \\ &= \frac{1}{2(1 - \tanh(2\beta))} \log(n_s) + O(\log(\log(n_s))). \end{aligned}$$

The above results hold for continuous time Glauber dynamics. However, we use discrete time Glauber dynamics, and thus adopt a modified version of these results. Since in the continuous case, there are in total n_s^d rate-one Poisson clocks, in the discrete case, it is natural to expect a slowdown of n_s^d times in the spectral gap, and mixing time. Therefore, we write

$$\hat{\gamma}^{1,Gl} := \frac{1 - \tanh(2\beta)}{n_s}, \quad \hat{t}_{\text{mix}}^{1,Gl} := \frac{1}{2(1 - \tanh(2\beta))} n_s \log(n_s). \quad (5.11)$$

In the 2-dimensional case, although (5.10) still applies for the continuous time Glauber dynamics, the explicit form of $\gamma_\infty^{d,Gl}$ as a function of β and h is not known. Therefore, we will use a modified version of (5.9):

$$\hat{\gamma}^{2,Gl} := \frac{1 - \beta/\beta_c}{n_s^2}, \quad \hat{t}_{mix}^{2,Gl} := \frac{1}{2} \frac{n_s^2 \log((1 - \beta/\beta_c)^2 n_s^2)}{1 - \beta/\beta_c}, \quad (5.12)$$

where $\beta_c = \frac{1}{2} \log(1 + \sqrt{2})$ is the critical inverse-temperature.

5.1.3. Simulation results for total magnetization

We will be interested in the total magnetization, i.e.

$$f(\sigma) = m(\sigma) := \sum_{i=1}^{n_s} \sigma_i.$$

In the case of no outside field ($h = 0$), we have, by symmetry, $\mathbb{E}m(\sigma) = 0$, for all of our models.

For some fixed β, h , and random initial distribution (uniformly chosen in σ), we run m Markov chains. The simulation results are shown in Figure 1. We observe that in our examples, the Bernstein-inequality, based on (4.9), provides a tight upper bound on the tail probabilities. In contrast, the normal quantiles based on the monotone sequence estimator of $\hat{\sigma}^2$ commonly underestimate the number of samples needed for a certain estimate precision. The Hoeffding-inequality does not incorporate the variance of f , and thus gives a weak bound on the tail probabilities.

For the Ricci curvature method, we plot (3.16). We choose the distance $d(x, y)$ as the Hamming distance, i.e. $d(x, y) = \sum_{i \leq n_s} \mathbb{1}[x_i \neq y_i]$. Then $\|f\|_{\text{Lip}} = 2$. In the case of the Ising models, Example 17 of Ollivier (2009) shows that

$$\kappa \geq \frac{1}{n_s} \left(1 - v_{\max} \frac{e^\beta - e^{-\beta}}{e^\beta + e^{-\beta}} \right), \quad (5.13)$$

with $v_{\max}$ being the maximum number of neighbors in the graph ($v_{\max}^{1D} = 2, v_{\max}^{2D} = 4$). For the Curie-Weiss model, because (5.7) contains $1/n_s$ in the sum, the bound becomes

$$\kappa_{CW} \geq \frac{1}{n_s} \left(1 - (n_s - 1) \frac{e^{\beta/n_s} - e^{-\beta/n_s}}{e^{\beta/n_s} + e^{-\beta/n_s}} \right). \quad (5.14)$$

Furthermore, in every case, we have $n_x \geq 1$, $\sigma_\infty \leq n_s$, and $\sigma^2(x) \leq 2$, $E(x) \leq n_s$. The bias term (3.17) is negligibly small with our parameters.




Fig 1: Simulation results for lattice models. The simulation result is plotted according to (5.5). When formulas are available for the mixing time and the spectral gap ((a),(c) and (e), see Section 5.1.2), the Hoeffding bound and Bernstein bound are plotted according to (2.1) and (2.7) respectively. We use estimated values of the parameters $\hat{\gamma}_{\text{dat}}$, $\hat{t}_{\text{mix,dat}}$, $\hat{\sigma}^2$ and $\hat{V}_f$ (see Section 4), and plot the estimated Bernstein bound according to (4.9). We also show the quantiles of $N(0, \hat{\sigma}^2)$ arising from the CLT ($\hat{\sigma}^2$ is estimated as in Geyer (1992), see Section 3). The bound based on Ricci curvature method is plotted according to (3.16).

5.2. Bayesian model averaging

In this section we look at simulation results on the space of directed acyclic graphs (DAGs) for Bayesian model averaging.

5.2.1. Definition of the model

DAGs are commonly used to encode the factored representation of a high-dimensional joint probability distribution. Let us consider a graph $G = (\mathcal{X}, E)$, where $\mathcal{X} = (X_1, X_2, \dots, X_n)$, is a set of vertices, each representing a random variable, and E is a set of directed edges between these variables. For a node $X_i \in \mathcal{X}$, we denote the set of its parents as $Pa(X_i)$, where $X_j \in Pa(X_i)$ if and only if $(X_j, X_i) \in E$. The non-descendents of a node $Nd(X_i)$ consist of the nodes to which there is no directed path from X_i . The DAG structure entails conditional independence relations among the variables of the following form:

$$X_i \perp\!\!\!\perp Nd(X_i) \mid Pa(X_i), \quad 1 \leq i \leq n \quad (5.15)$$

With these independence assumptions, the joint distribution of the variables factors as

$$\mathbb{P}(X_1, X_2, \dots, X_n) = \prod_{i=1}^n \mathbb{P}(X_i \mid Pa(X_i)) \quad (5.16)$$

Now, given a set of observations D , we attempt to make predictions about a function f of the model structure G . D consists of vectors of realizations of $(X_1, X_2, \dots, X_n)$. One could find a single best DAG structure with respect to D , and use it to calculate the value of f . Instead, we follow a Bayesian model averaging approach, where we calculate the posterior probability of each possible DAG structure and use it as a weight when making the prediction. We refer to [Neapolitan \(2004\)](#) for a more detailed account of this approach. The prediction $\mathbb{E}[f(G) \mid D]$ can be expressed as a weighted average of individual predictions based on each possible DAG structure g :

$$\mathbb{E}[f(G) \mid D] = \sum_{g \in \Omega_n} f(g) \mathbb{P}(G = g \mid D), \quad (5.17)$$

where Ω_n is the set of all DAG structures on n variables.

The model G is parametrized using a set of conditional probability tables describing the probability of each node taking a certain value given its parents. We denote the set of all such parameters θ_G .

The posterior probability of a structure G can be obtained by applying Bayes theorem on its marginal likelihood. The marginal likelihood is generally expressed as

$$\mathbb{P}(D \mid G) = \int_{\theta_G} \mathbb{P}(D \mid \theta_G, G) \mathbb{P}(\theta_G) d\theta_G. \quad (5.18)$$

In our current example, we assume that each random variable is binary, that is, $X_i \in \{0, 1\}$. As is typically done in the context of binary DAG models, we set a beta distribution as the prior distribution of each variable conditioned on its parent configuration.

Using beta priors, [Heckerman, Geiger and Chickering \(1995\)](#) shows that the marginal likelihood can be calculated as

$$\mathbb{P}(D|G) = \prod_{i=1}^n \prod_{j=1}^{q_i} \frac{\Gamma(s_{ij})}{\Gamma(d_{ij} + s_{ij})} \prod_{k=1}^{r_i} \frac{\Gamma(d_{ijk} + s_{ijk})}{\Gamma(s_{ijk})}, \quad (5.19)$$

where i refers to a node X_i , j is a value configuration of the parents of node X_i , with q_i the total number of parent value configurations, k indicates the value of node X_i under parent configuration j , and r_i is the number of different values that X_i can take. For each combination of indices, d_{ij} and d_{ijk} represent the observed count, while s_{ij} and s_{ijk} are the prior counts. To make priors consistent among different DAG structures, we choose a fix equivalent sample size S , and set $s_{ijk} = \frac{S}{q_i r_i}$.

For simplicity, we assume that the prior probability for each structure is equal, that is, $\forall G \in \Omega_n, \mathbb{P}(G) = \frac{1}{|\Omega_n|}$.

As the number of summation terms in (5.17) can be prohibitively large to compute exactly, we design a Markov chain with stationary distribution $\mathbb{P}(G|D)$ and use a Monte Carlo estimate to approximate the prediction.

5.2.2. Procedure

We follow [Madigan, York and Allard \(1995\)](#), and design a Markov chain on Ω_n with stationary distribution $\mathbb{P}(G|D)$. Starting with an initial DAG structure, the chain either adds or removes a single edge at each proposal step. We denote the neighborhood of a state G_i as $Nb(G_i)$, which is the set of DAGs that differ from G_i by one edge addition or one edge removal. The chain then uses the following probabilities to propose the next state:

$$T(G_j|G_i) = \begin{cases} \frac{1}{|Nb(G_i)|}, & G_j \in Nb(G_i) \\ 0, & G_j \notin Nb(G_i) \end{cases}. \quad (5.20)$$

When making the proposal, we make sure that only valid (cycle-free) DAGs are considered. The chain moves to the proposed state with the following acceptance probability:

$$A(G_j|G_i) = \min \left\{ 1, \frac{|Nb(G_i)| \mathbb{P}(G_j|D)}{|Nb(G_j)| \mathbb{P}(G_i|D)} \right\}. \quad (5.21)$$

Note: The ratio of marginal likelihoods can be evaluated locally at the target node of the single edge that is changed during the proposal step. As opposed to some of the lattice models discussed in the previous section, here, no analytic formulas are known for the mixing time and spectral gap of the Markov chain.

5.2.3. Simulation results

In the following simulation example, we have a set of $n = 6$ variables, thus the space of the Markov Chain consist of DAGs with 6 vertices. We take a data set D consisting of 20 vectors generated from a known DAG on 6 nodes (structure not shown), and assume a prior equivalent




Fig 2: Simulation results for Bayesian model averaging. The simulation result is plotted according to (5.5). We use estimated values of the parameters $\hat{\gamma}_{dat}$, $\hat{t}_{mix,dat}$, $\hat{\sigma}^2$ and $\hat{V}_f$ (see Section 4), and plot the estimated Bernstein-bound according to (4.9). We also show the quantiles of $N(0, \hat{\sigma}^2)$, arising from the CLT ($\hat{\sigma}^2$ is estimated as in Geyer (1992), see Section 3).

sample size of 4. Our goal is to estimate the posterior probability of an edge being present in the structure:

$$f(G) = \begin{cases} 1, & (X_i, X_j) \in E_G \\ 0, & (X_i, X_j) \notin E_G \end{cases}. \quad (5.22)$$

We look at two cases, first, at the presence of the edge $e_a = (i = 1, j = 2)$, and then at $e_b = (i = 1, j = 4)$. The simulation results are shown in Figure 2 (a) and (b) respectively. These figures show examples of exponential tails, for which our proposed Bernstein bound provides a tight upper bound. The normal quantile based estimate is poor on the side with exponential tail.

6. Final remarks

In order to get rigorous, sharp error bounds for empirical averages in MCMC, one needs to know the mixing time of the chain (for setting the “burn-in time” t_0 sufficiently large), the spectral gap (for reversible chains), and the concentration properties of the function f at the stationary distribution. The Hoeffding inequalities only use the lower and upper bounds on f , while Bernstein inequalities take into account the variance of f as well. Our simulation results show that this distinction is important for obtaining tight error bounds. While the normal approximation by the central limit theorem can only handle Gaussian tails, concentration inequalities are also applicable in case of exponential tails that arise in practice.

It would be interesting to get even sharper results, under additional conditions on f . For instance if f has Gaussian or exponential tails (such tail inequalities are proven for statistical physical systems satisfying the Dobrushin condition in Paulin (2012b)), one could get sharper error bounds, since it is the typical deviation of f that really matters and not its maximal range.

For further practical examples where the bounds we have presented can be used, we refer the reader to [Gilks, Richardson and Spiegelhalter \(1995\)](#), [Liu \(2008\)](#) and [Landau and Binder \(2009\)](#).

Acknowledgements

The second author thanks Doma Szász and Mogyi Tóth for infecting him with their enthusiasm of probability. He also thanks his brother, Roland Paulin, for the enlightening discussions. The authors thank their thesis supervisors, Louis Chen, Adrian Röllin and David Hsu for the opportunity to study in Singapore, and their useful advices. We also thank Daniel Rudolf for his comments. Finally, we thank Lee Hwee Kuan for his contribution to the simulation code.

References

- BROOKS, S. P. and GELMAN, A. (1998). General methods for monitoring convergence of iterative simulations. *J. Comput. Graph. Statist.* **7** 434–455. . [MR1665662 \(99k:62055\)](#)
- BUBLEY, R., DYER, M. et al. (1997). Path coupling, Dobrushin uniqueness, and approximate counting. *Research Report Series - University of Leeds School of Computer Studies LU SCS RR*. Available at http://reference.kfupm.edu.sa/content/p/a/path_coupling_dobrushin_uniqueness_and_9577
- CHAZOTTES, J. R., COLLET, P., KÜLSKE, C. and REDIG, F. (2007). Concentration inequalities for random fields via coupling. *Probab. Theory Related Fields* **137** 201–225. . [MR2278456 \(2008i:60167\)](#)
- COWLES, M. K. and CARLIN, B. P. (1996). Markov chain Monte Carlo convergence diagnostics: a comparative review. *J. Amer. Statist. Assoc.* **91** 883–904. . [MR1395755](#)
- DIACONIS, P. (2011). The mathematics of mixing things up. *J. Stat. Phys.* **144** 445–458. . [MR2826629 \(2012j:60207\)](#)
- DIACONIS, P., HOLMES, S. and NEAL, R. M. (2000). Analysis of a nonreversible Markov chain sampler. *Ann. Appl. Probab.* **10** 726–752. . [MR1789978 \(2001i:60114\)](#)
- DING, J., LUBETZKY, E. and PERES, Y. (2009). The mixing time evolution of Glauber dynamics for the mean-field Ising model. *Comm. Math. Phys.* **289** 725–764. . [MR2506768 \(2010e:82064\)](#)
- GELMAN, A. and RUBIN, D. B. (1992). Inference from iterative simulation using multiple sequences. *Statistical science* **7** 457–472.
- GELMAN, A., CARLIN, J. B., STERN, H. S. and RUBIN, D. B. (2004). *Bayesian data analysis*, second ed. *Texts in Statistical Science Series*. Chapman & Hall/CRC, Boca Raton, FL. [MR2027492 \(2004j:62001\)](#)
- GEYER, C. J. (1992). Practical Markov chain Monte Carlo. *Statistical Science* **7** 473–483.
- GILKS, W. R., RICHARDSON, S. and SPIEGELHALTER, D. (1995). *Markov chain Monte Carlo in practice: interdisciplinary statistics* **2**. Chapman & Hall/CRC.
- HECKERMAN, D., GEIGER, D. and CHICKERING, D. M. (1995). Learning Bayesian networks: The combination of knowledge and statistical data. *Machine learning* **20** 197–243.
- JANSON, S. (2004). Large deviations for sums of partly dependent random variables. *Random Structures Algorithms* **24** 234–248. . [MR2068873 \(2005e:60061\)](#)

- JOULIN, A. and OLLIVIER, Y. (2010). Curvature, concentration and error estimates for Markov chain Monte Carlo. *Ann. Probab.* **38** 2418–2442. . [MR2683634 \(2011j:60229\)](#)
- KIPNIS, C. and VARADHAN, S. R. S. (1986). Central limit theorem for additive functionals of reversible Markov processes and applications to simple exclusions. *Comm. Math. Phys.* **104** 1–19. [MR834478 \(87i:60038\)](#)
- KONTOROVICH, L. (2007). *Measure Concentration of Strongly Mixing Processes with Applications*. Ph.D. dissertation, Carnegie Mellon University, Available at <http://www.cs.bgu.ac.il/~karyeh/thesis.pdf>.
- LANDAU, D. P. and BINDER, K. (2009). *A guide to Monte Carlo simulations in statistical physics*, Third ed. Cambridge University Press, Cambridge. . [MR2559932 \(2011a:82046\)](#)
- LATUSZYSKI, K., MIAOJEDOW, B. and NIEMIRO, W. (2011). Nonasymptotic bounds on the estimation error of MCMC algorithms. *arXiv preprint arXiv:1106.4739*.
- LEÓN, C. A. and PERRON, F. (2004). Optimal Hoeffding bounds for discrete reversible Markov chains. *Ann. Appl. Probab.* **14** 958–970. . [MR2052909 \(2005d:60109\)](#)
- LEVIN, D. A., PERES, Y. and WILMER, E. L. (2009). *Markov chains and mixing times*. American Mathematical Society, Providence, RI. With a chapter by James G. Propp and David B. Wilson. [MR2466937 \(2010c:60209\)](#)
- LEZAUD, P. (1998a). Chernoff-type bound for finite Markov chains. *Ann. Appl. Probab.* **8** 849–867. . [MR1627795 \(99f:60061\)](#)
- LEZAUD, P. (1998b). *Etude quantitative des chaînes de Markov par perturbation de leur noyau*. Thèse doctorat mathématiques appliquées de l'Université Paul Sabatier de Toulouse, Available at http://pom.tls.cena.fr/papers/thesis/these_lezaud.pdf.
- LIU, J. S. (2008). *Monte Carlo strategies in scientific computing*. Springer Series in Statistics. Springer, New York. [MR2401592 \(2010b:65013\)](#)
- LUBETZKY, E. and SLY, A. (2009). Cutoff for the Ising model on the lattice. *Inventiones Mathematicae* 1–37.
- MADIGAN, D., YORK, J. and ALLARD, D. (1995). Bayesian graphical models for discrete data. *International Statistical Review/Revue Internationale de Statistique* 215–232.
- MARTON, K. (1996). Bounding $\bar{d}$ -distance by informational divergence: a method to prove measure concentration. *Ann. Probab.* **24** 857–866. . [MR1404531 \(97f:60064\)](#)
- METROPOLIS, N. (1987). The beginning of the Monte Carlo method. *Los Alamos Sci.* 15, Special Issue 125–130. Stanislaw Ulam 1909–1984. [MR935771](#)
- MEYN, S. and TWEEDIE, R. L. (2009). *Markov chains and stochastic stability*, Second ed. Cambridge University Press, Cambridge. With a prologue by Peter W. Glynn. [MR2509253 \(2010h:60206\)](#)
- NEAPOLITAN, R. E. (2004). *Learning Bayesian networks*. Pearson Prentice Hall Upper Saddle River, NJ.
- OLLIVIER, Y. (2009). Ricci curvature of Markov chains on metric spaces. *J. Funct. Anal.* **256** 810–864. . [MR2484937 \(2010j:58081\)](#)
- PAULIN, D. (2012a). Concentration inequalities for Markov chains by Marton couplings. *arXiv preprint*.
- PAULIN, D. (2012b). Concentration of Self-Bounding Functions in Weakly Dependent Spaces by Stein's Method. *arXiv preprint*.
- RUDOLF, D. (2011). Explicit error bounds for Markov chain Monte Carlo. *arXiv preprint arXiv:1108.3201*.

7. Appendix

Proof of Theorem 2.3. Marton (1996) proves measure concentration in Hamming distance for countable state Markov chains. For a homogeneous, ergodic Markov chain with state space Ω , and transition probabilities $P_{i,j}$, let us denote

$$q := \max_{i,j \in \Omega} d_{TV}(P_{i,\cdot}, P_{j,\cdot}). \quad (7.1)$$

Then Proposition 1 of Marton (1996) proves that measure concentration holds with constants $1/(1-q)^2$ times worse than in the independent case (see also Kontorovich (2007) and Chazottes et al. (2007)). In particular, Mcdiarmid's bounded differences inequality holds with $1/(1-q)^2$ times weaker constant than in the independent case:

Proposition 7.1. *Suppose that $g : \Omega^n \rightarrow \mathbb{R}$ is C -Hamming Lipschitz (i.e. $g(x)$ can change at most by C if we change only one coordinate in x), and let $X = (X_1, \dots, X_n)$ be a homogeneous, ergodic Markov chain taking values in Ω , then for every λ ,*

$$\log \mathbb{E} \exp(\lambda(g(X) - \mathbb{E}g(X))) \leq \frac{\lambda^2 C n}{8(1-q)^2}, \quad (7.2)$$

and thus

$$\mathbb{P}(g(X) \geq \mathbb{E}g(X) + t), \mathbb{P}(g(X) \leq \mathbb{E}g(X) - t) \leq \exp\left(-\frac{2(1-q)^2 t^2}{C n}\right), \quad (7.3)$$

Fix some $0 \leq \epsilon < 1/2$. Suppose, without loss of generality, that N is divisible by $t_{\text{mix}}(\epsilon)$, and let $n = N/t_{\text{mix}}(\epsilon)$. Divide $X_1, \dots, X_N$ into $t_{\text{mix}}(\epsilon)$ groups such that the indexes of the elements in these groups are at least $t_{\text{mix}}(\epsilon)$ distance from each other:

$$\begin{aligned} Y^{(1)} &:= (Y_1^{(1)}, \dots, Y_n^{(1)}) := (X_1, X_{1+t_{\text{mix}}(\epsilon)}, \dots, X_{1+(n-1)t_{\text{mix}}(\epsilon)}), \\ &\vdots \\ Y^{(n)} &:= (Y_1^{t_{\text{mix}}(\epsilon)}, \dots, Y_n^{t_{\text{mix}}(\epsilon)}) := (X_{t_{\text{mix}}(\epsilon)}, X_{2t_{\text{mix}}(\epsilon)}, \dots, X_{nt_{\text{mix}}(\epsilon)}). \end{aligned}$$

Now we use a trick from the proof of Theorem 1 of Janson (2004). Denote

$$W := \sum_{i=1}^N f(X_i) - \mathbb{E}f(X_i) = \sum_{j=1}^{t_{\text{mix}}(\epsilon)} \sum_{i=1}^n f(X_{t_{\text{mix}}(\epsilon)(i-1)+j}) - \mathbb{E}f(X_{t_{\text{mix}}(\epsilon)(i-1)+j}),$$

then by Jensen's inequality,

$$\begin{aligned} \mathbb{E}(\exp(\lambda W)) &\leq \frac{1}{t_{\text{mix}}(\epsilon)} \cdot \\ &\cdot \sum_{j=1}^{t_{\text{mix}}(\epsilon)} \mathbb{E} \left(\exp \left(t_{\text{mix}}(\epsilon) \cdot \lambda \sum_{i=1}^n [f(X_{t_{\text{mix}}(\epsilon)(i-1)+j}) - \mathbb{E}f(X_{t_{\text{mix}}(\epsilon)(i-1)+j})] \right) \right). \end{aligned}$$

Now we notice that $\{X_{t_{\text{mix}}(\epsilon)(i-1)+j}\}_{1 \leq i \leq n}$ is a Markov chain by itself, and it is easy to see that for this chain, $q \leq 2\epsilon$, thus we can apply (7.2), with $C = b - a$:

$$\mathbb{E}(\exp(\lambda W)) \leq \exp \left(\frac{\lambda^2 n(b-a) \cdot t_{\text{mix}}}{8(1-2\epsilon)^2} \right),$$

and thus, by Markov's inequality,

$$\mathbb{P} \left(\sum_{i=1}^N f(X_i) \geq \sum_{i=1}^N \mathbb{E} f(X_i) + t \right) \leq \exp \left(\frac{-2t^2(1-2\epsilon)^2}{N(b-a)^2 t_{\text{mix}}(\epsilon)} \right).$$

To get (2.11), we only need to rescale this, change N to $N - t_0$, optimize in ϵ , and show that

$$\left| \frac{1}{N - t_0} \sum_{i=t_0+1}^N \mathbb{E} f(X_i) - \mathbb{E}_\pi f \right| \leq \frac{\eta(t_0)(b-a)}{N - t_0},$$

these are left to the reader. $\square$

Proof of Proposition 4.1. We have

$$\begin{aligned} V_f &= \mathbb{E}_\pi f^2 - (\mathbb{E}_\pi f)^2, \text{ and} \\ \hat{V}_f &= \left(\frac{1}{N - t_0} \sum_{i=t_0+1}^N f^2(X_i) \right) - \left(\frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0} \right)^2. \end{aligned}$$

Define

$$\begin{aligned} D_1 &:= \mathbb{E}_\pi f^2 - \frac{1}{N - t_0} \sum_{i=t_0+1}^N f^2(X_i), \text{ and} \\ D_2 &:= \left(\frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0} \right)^2 - (\mathbb{E}_\pi f)^2, \text{ then} \\ V_f - \hat{V}_f &= D_1 + D_2. \end{aligned}$$

The upper tail of D_1 can be bounded by Theorem 2.3:

$$\mathbb{P}(D_1 \geq \eta_{\min}(t_0)C^2 + t) \leq \exp \left(-\frac{2t^2(N - t_0)}{C^4 t_{\min}^{\min}} \right). \quad (7.4)$$

Now we bound the upper tail of D_2 :

$$\begin{aligned} D_2 &= \left(\frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0} \right)^2 - (\mathbb{E}_\pi f)^2 \\ &= \left(\frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0} - \mathbb{E}_\pi f \right) \left(\frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0} + \mathbb{E}_\pi f \right) \\ &= \left(\frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0} - \mathbb{E}_\pi f \right) \left(2 \frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0} + \mathbb{E}_\pi f - \frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0} \right) \\ &\leq \left(\frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0} - \mathbb{E}_\pi f \right) \left(2 \frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0} \right) \\ &\leq 2C \left| \frac{\sum_{i=t_0+1}^N f(X_i)}{N - t_0} - \mathbb{E}_\pi f \right|, \end{aligned}$$

therefore Theorem 2.3 gives

$$\mathbb{P}(D_2 \geq 4\eta_{\min}(t_0)C^2 + t) \leq 2 \exp\left(-\frac{2t^2(N - t_0)}{4C^4 t_{\min}^{\min}}\right). \quad (7.5)$$

Combining (7.4) (for $t/3$) and (7.5) (for $2t/3$), we get

$$\begin{aligned} \mathbb{P}(D_1 + D_2 \geq 5\eta_{\min}(t_0)C^2 + t) &\leq \exp\left(-\frac{2(t/3)^2(N - t_0)}{C^4 t_{\min}^{\min}}\right) + \\ &2 \exp\left(-\frac{2(2/3t)^2(N - t_0)}{4C^4 t_{\min}^{\min}}\right), \end{aligned}$$

so (4.3) follows. The proof of (4.4) is similar. □